

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: A61K 49/00, 31/54, C07D 279/12	A1	(11) International Publication Number: WO 97/26018 (43) International Publication Date: 24 July 1997 (24.07.97)
(21) International Application Number: PCT/US96/00509 (22) International Filing Date: 16 January 1996 (16.01.96) (71) Applicant (for all designated States except US): ZILA PHARMACEUTICALS, INC. [US/US]; 5227 North 7th Street, Phoenix, AZ 85014-2800 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): POMERANTZ, Edwin [US/US]; 22662 Cavalier Street, Woodland Hills, CA 91364 (US). (74) Agent: DRUMMOND, William, H.; Drummond & Duckworth, Suite 500, 4590 MacArthur Boulevard, Newport Beach, CA 92660 (US).		(81) Designated States: AU, CA, JP, KR, RU, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: METHODS AND COMPOSITIONS FOR IN-VIVO DETECTION OF ORAL CANCERS AND PRECANCEROUS CONDITIONS (57) Abstract <p>A method for in-vivo detection of premalignant oral lesions and oral carcinomas includes the step of applying to oral tissue a non-toxic dye other than toluidine blue O, selected from compounds having structure (I), and ionic derivatives thereof, wherein R¹ is H, or lower alkyl group or N(R³)₂, R² is N, S or O, and R³ is H or a lower alkyl group.</p> <div data-bbox="992 1150 1344 1312"><p style="text-align: right;">(I)</p></div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

-1-

METHODS AND COMPOSITIONS FOR IN-VIVO DETECTION
OF ORAL CANCERS AND PRECANCEROUS CONDITIONS

5 This invention relates to in vivo methods for
detection of premalignant oral lesions and oral
carcinomas.

In another respect the invention pertains to
compositions for carrying out such diagnostic procedures.

10 In another aspect the invention relates to such in
vivo diagnostic procedures and compositions useful
therein which are especially useful in screening patients
for possible oral cancer as part of routine dentist's or
physician's examinations, or procedures such as periodic
check-up's, dental cleaning, etc.

15 In yet another aspect the invention relates to such
procedures and compositions which utilize dye stains
which are more readily available and/or less expensive
and less complicated to synthesize and/or purify than the
dyes employed in prior art procedures.

20 In yet another respect the invention concerns such
in vivo procedures and compositions employing dyes which
are less potentially toxic than prior art fluorescein
derivatives and methylene blue and which may, therefore,
be applied by rinsing the entire oral cavity and/or
gargling.

-2-

In-vivo diagnostic procedures for detection of premalignant oral lesions or oral carcinomas, employing dye compositions which are selectively retained by tissues rendered abnormal due to dysplasia, hyperplasia, tumorigenesis, and other active surface lesions, are known in the art. For example, procedures employing fluorescein or fluorescein derivatives are disclosed in Chenz, Chinese Journal of Stomatology (27:44-47 (1992)) and Filiurin (Stomatologiia (Russian) 72:44-47 (1993)). These procedures involve application of the dye, followed by examination under ultraviolet light to detect the cancerous/precancerous tissue, which is selectively fluorescent.

Another prior art procedure involves in-vivo application by rinsing with toluidine blue O, followed by normal visual examination to detect any selectively stained tissue. Such procedures are disclosed, for example in the U.S. Patent to Tucci, et al. No. 5,372,801 and in U.S. Patent 4,321,251 to Mashberg. Toluidine blue, has been used for decades as a histopathological stain for in-vitro use. Through this use it has become known as a metachromatic dye, staining nuclei rich in DNA and RNA a purple to pink color. The inherent deep blue color of toluidine blue is changed to purple or pink when the dye is bound to a nucleic acid or other acidic cellular macromolecule. Of course, this type of staining is dependent on the dye gaining access to internal

-3-

subcellular structures such as the nucleus. Such access is readily obtained only by "fixing" a tissue sample of formaldehyde or other reagent that disrupts the cellular membrane without destroying general cellular structure.

5 In contrast to the mechanism involved with in-vitro use, the staining of oral tissue in-vivo by toluidine blue O is due to its ability to penetrate the enlarged and more accessible interstitial spaces of dysplastic tissue, which retains the dye longer than areas
10 containing normal tissue with its tight intercellular junctions. Empirically, the observation is made that toluidine blue simply does not penetrate normal intercellular spaces efficiently than those of normal tissue, and becomes temporarily localized, it diffuses
15 out with time and does not remain bound to the cells as it would if it actually "stained" acidic substructures.

 The Mashberg procedure involves application of the toluidine blue O solution as a rinse of the entire oral cavity, with gargling, following by rinses with water and
20 acidic acid to remove the dye which is not retained by the cancerous or precancerous tissue. The preliminary Mashberg diagnosis is then confirmed by direct application of the toluidine blue O composition to the suspect site 10-14 days later.

-4-


The Tucci '801 patent discloses an improved toluidine blue O composition for use according to the general Mashberg procedure.

5 An in-vivo procedure involving use of Lugol's solution (iodine) and toluidine blue O was proposed for detecting esophageal cancer synchronous with upper aerodigestive tract cancers in Papazian (Gastroenterologie Clinique et Biologique (9:16-22, 1985). Because of possible toxic properties of
10 fluorescein, iodine and methylene blue, it is considered that they are not suitable for indiscriminate application by oral rinse and/or gargling procedures. Toluidine blue O is not toxic, but has been expensive to synthesize and purify in commercial quantities.

15 Accordingly, it would be advantageous to employ dyes other than fluorescein, fluorescein derivatives, iodine, methylene blue or toluidine blue O in diagnostic procedures for in-vivo detection of premalignant oral lesions and oral carcinomas.

20 I have now discovered that non-toxic dyes other than methylene blue or toluidine blue O, selected from compounds having the structure

-5-



and ionic derivatives thereof, wherein R¹ is H, a lower alkyl group or N(R³)₂, R² is N, S or O, and R³ is H or a lower alkyl group, are effective for in-vivo detection of premalignant oral lesions or oral carcinomas. The lower alkyl groups can be C₁-C₄, straight or branched chain groups, preferably methyl or ethyl groups. In general, these dyes are employed according to the same general procedure as disclosed by the Mashberg '251 patent, i.e., by sequentially rinsing the oral cavity with the dye stain composition, which is selectively retained by cancerous and precancerous tissues, and then rinsing the oral cavity to remove unretained stain composition.

The determination of whether a particular dye having the above-described structure is selectively retained can be readily ascertained by persons skilled in the art without undue experimentation, by using the dye in accordance with the general Mashberg procedure, described above. The potential toxicity of a particular dye can be readily determined by art-recognized techniques and standards, e.g., animal feeding studies.

-6-

Suitable compositions for the application of the dyes defined above are prepared by mixing the dye with a pharmaceutically acceptable oxidizing agent which oxidizes any leuco form of the dye to the chromo form.

5 The oxidizing agents are selected to be pharmaceutically acceptable, i.e., are non-toxic and do not cause undesirable side reactions which degrade the dye. Those skilled in the art will be able to identify suitable oxidizing agents for use in accordance with the invention

10 by routine tests of known non-toxic mild oxidants. For example, in accordance with the presently preferred embodiment of the invention it is preferred to use hydrogen peroxide, either directly or by reaction of known peroxide precursors with water in the compositions,

15 e.g., water soluble "per" compounds, such as urea peroxide,, sodium perborate tetrahydrate, sodium percarbonate, calcium peroxide and the like.

The compositions of the invention are preferably formulated to yield a final solution which is

20 substantially isotonic and has a pH in the range of approximately 2.5 to 7.0, preferably 4.0 to 5.0. This is accomplished by adding an appropriate water soluble buffer system. The presently preferred buffer system is acetic acid-sodium acetate. Other suitable buffer

25 systems include citric acid-sodium citrate, or mixed acid-salt systems such as citric acid-sodium phosphate and the like.

-7-

The solvent used to provide liquid dye compositions is an aqueous solvent. According to the preferred embodiment of the invention, the solvent includes a pharmaceutically acceptable, i.e., non-toxic, non-
5 reactive, alcohol, e.g., ethyl alcohol. Such solvents do not appreciably interfere with the tissue staining mechanism and do not themselves contribute to the reduction of chromo forms of the dyes to the leuco forms.

Flavoring, stable to the oxidizing agent, may be
10 added to improve the palatability of the dye solution if it is to be used as a "rinse".

The amount of the dye in the liquid compositions is preferably adjusted to yield a concentration of approximately 1% by weight in the final composition,
15 although higher concentrations can be employed and lower concentrations are at least partially effective, e.g., from about 0.5 to about 3.5 weight %.

Surprisingly, dyes having the above-defined structure, which are closely related to toluidine blue O,
20 e.g., methylene blue and thionine, have proven to be ineffective, whereas other dyes closely related to toluidine blue O, e.g., Brilliant Cresyl Blue, Azure A, Azure B and Azure C are effective.

- 8 -

According to my present knowledge, Azure B and Azure C are employed in accordance with the presently preferred embodiment of the invention.

5 The invention also contemplates compositions for use in accordance with the methods of the invention and processes for manufacturing such composition, in which any leuco form of the dye present in the composition is oxidized to the chromo form, by inclusion of a pharmaceutically acceptable oxidizing agent in the
10 composition, in the manner analogous to that disclosed in the Tucci et al. USA Patent 5,372,801.

EXAMPLES

The following examples are presented in order to illustrate practice of the invention to those skilled in
15 the art and not by way of limitation on the scope thereof.

EXAMPLE 1

Test compositions of Azure B, Azure C, Brilliant Cresyl Blue ("BCB") and thionine dyes and a control
20 composition, toluidine blue O dye ("TBO") are prepared by mixing each of the dyes with the indicated ingredients in the following proportions (% by weight):

-9-

	<u>Ingredient</u>	<u>% by Weight</u>
	Purified Water, U.S.P.	83.86
	Glacial Acetic Acid, U.S.P.	4.61
	Sodium Acetate Trihydrate, U.S.P.	2.45
5	SD18 Ethyl Alcohol	7.48
	Hydrogen Peroxide 30%, U.S.P.	0.41
	IFF Raspberry IC563457	0.20
	Dye	1.00

EXAMPLE 2

10 A rinse solution is prepared by mixing the following ingredients in the indicated proportions (weight %):

	Purified Water, U.S.P.	98.70
	Glacial Acetic Acid, U.S.P.	1.00
	Sodium Benzoate, U.S.P.	0.10
15	IFF Raspberry IC563457	0.20

EXAMPLE 3

The clinical effectiveness of each of the test mixtures, compared to the TBO control mixture, is determined, utilizing the Mashberg protocol.

20 Patients are first screened for oral pathology employing the TBO control mixture. After identifying potential cancerous or precancerous pathology, all traces

-10-

of the TBO are removed by repeated rinses with water and the acetic acid rinse solution. Those patients exhibiting oral pathology are then used as test subjects for each of the test mixtures. 2-3 cc of each test mixture is applied by painting the pathologic mucosal surfaces, followed by rinsing with the rinse mixture and water to remove excess dye composition. After examination with each dye and before testing with the next dye, all traces of the dye are removed from the lesion by repeated rinsing with the rinse mixture and water.

The Azure B test mixture provides equivalent staining intensity to TBO, localization of the lesions and delineation of the borders of the lesions. Adjacent unstained tissue shows slight dye accumulation, but not to the detriment of lesion discrimination.

The Azure C test mixture provides equivalent staining intensity to TBO, localization of the lesions and delineation of the borders of the lesions. Adjacent, uninvolved tissues are stained to a red/purple hue, but not to the detriment of lesion discrimination.

The Brilliant Cresyl Blue test mixture provides results similar to Azure B and Azure C.

-11-

The Thionine test mixture provides a warm red background stain on both the stained and unstained tissues, such that the discrimination of the lesion borders is insufficient.

5

EXAMPLE 4

The procedures of Example 3 are repeated, except that the test and control compositions are applied to the oral mucosa by rinsing, with gargling, instead of by direct application to the locus of the suspect sites.

10 Equivalent results are obtained.

Having disclosed my invention in such terms as to enable those skilled in the art to make and use it, and, having identified the presently preferred embodiments thereof, I CLAIM:


-12-

1. In a method for in vivo detection of premalignant oral lesions and oral carcinomas, including the steps of sequentially rinsing the oral cavity with

a dye stain composition which is selectively retained by cancerous and precancerous tissues, and

a rinse composition for removing unretained stain composition,

the step of applying to oral tissue, a stain composition comprising a non-toxic dye other than toluidine blue O and methylene blue, selected from compounds having the structure



and ionic derivatives thereof, wherein R^1 is H, a lower alkyl group or $N(R^3)_2$, R^2 is N, S or O, and R^3 is H or a lower alkyl group.

-13-

2. A method for in vivo detection of premalignant oral lesions and oral carcinomas, including the steps of sequentially rinsing the oral cavity with

5 a dye stain composition which is selectively retained by cancerous and precancerous tissues, and

a rinse composition for removing unretained stain composition,

10 the step of applying to oral tissue, a stain composition comprising a dye selected from the group consisting of Azure B, Azure C and Brilliant Cresyl Blue.

3. The method of Claim 2 in which said dye is Brilliant Cresyl Blue.

15 4. The method of Claim 2 in which said dye is selected from the group consisting of Azure B and Azure C.


5. The method of Claim 4 in which said dye is Azure B.

6. The method of Claim 4 in which said dye is Azure C.

-14-

7. A biological stain composition for in vivo detection of cancerous and precancerous tissue, comprising:

- (a) a non-toxic dye other than toluidine blue O and methylene blue having the structure



and ionic derivatives thereof, wherein R^1 is H, a lower alkyl group or $(R^3)_2$, R^2 is N, S or O, and R^3 is H or a lower alkyl group, and

- (b) a pharmaceutically acceptable oxidizing agent for leuco forms of the dye.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/00509

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 49/00, 31/54; C07D 279/12

US CL :424/9.6; 514/227.5; 544/56

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/9.6, 7.1; 514/227.5, 222.2; 544/56, 2, 3

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, MEDLINE, BIOSIS, CAPLUS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 3,852,413 A (P.S. CAMMARATA) 03 December 1974, column 3, lines 13-40	1-7
Y	US 4,321,251 A (A. MASHBERG) 23 March 1982, columns 1 and 2, abstract.	1-7
Y	US 4,950,665 A (R.A. FLOYD) 21 August 1990, column 1, lines 47-50 and column 3, lines 36-45.	1-7
Y	US 5,372,801 A (M.K. MALMROS ET AL.) 13 December 1994, column 1, lines 17-22.	1-7



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

06 MAY 1996

Date of mailing of the international search report

15 MAY 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Michael G. Hartley

Telephone No. (703) 308-1235